

RECENT ADVANCES IN STUDIES OF HOMING MECHANISMS

LYNN L. ROGERS, USDA— Forest Service, North Central Forest Experiment Station, 1992 Folwell Avenue, St. Paul, MN 55108

The ability of certain species to orient homeward from unfamiliar areas has long been studied (Matthews 1968, Emien 1975). Three levels of navigation ability have been defined (Griffin 1955). Type I is simple orientation by means of familiar landmarks (visual or otherwise). Type II is the ability to move in a particular compass direction without reference to landmarks; and Type III, often called true navigation, is the ability to orient homeward (or toward some other goal) from a variety of unfamiliar areas. There is evidence that homing pigeons (Walcott 1974), purple martins (*Progne subis*) (Southern 1959, 1968), the Manx shearwater (*Puffinus puffinus*) (Mazzeo 1953), the Laysan albatross (*Diomedea immutabilis*) (Kenyon and Rice 1958), and black bears (*Ursus americanus*) (Harger 1970, Alt 1977, Rogers 1984, in preparation a and b) possess Type III navigational ability.

Study of homing mechanisms has concentrated on homing pigeons and has led to the discovery of several surprising sensory abilities in this species. They can detect polarized light (Delius et al. 1976, Kreithen 1978), ultraviolet light (Kreithen 1978, Delius and Emmerton 1978), very low frequency infrasounds (Kreithen 1978), and changes in atmospheric pressure (Delius and Emmerton 1978), but which of these senses might be used to determine the homeward direction is unknown.

Magnetite has been discovered in the heads of homing pigeons (Walcott et al. 1979), dolphins (Zoeger et al. 1981), and man (Baker et al. 1983) and was found in the bodies of bees (Gould et al. 1978) and certain bacteria (Frankel et al. 1979). A current area of concentrated study is determining which vertebrates possess magnetite and use it to orient to the earth's magnetic field. Only the bacteria are known to use magnetite for orientation (Frankel et al. 1979), but the diversity of groups that have magnetite has led to speculation that geomagnetic compass sense may be widespread (Baker et al. 1983). However, even if some species have this sense, the mechanisms of true navigation remain a mystery because a mental map is still needed to use a compass sense.

There is strong evidence that odors are important in the formation of the mental map in homing pigeons (Walraff 1983). Study has concentrated on the role of odors in navigation following reports by Papi et al. (1973) and Benvenuti et al. (1973a, 1973b) that homing pigeons whose sense of smell is experimentally impaired show poor homing success. Papi et al. (1973) proposed that "Pigeons, during their first months of life, would learn to recognize the odors prevailing in the area of the loft, but when the wind blows, they would also receive information about foreign odors from surrounding areas. At the same time they would

associate the different foreign odors with the direction from which they come." Followup experiments were conducted in which pigeons were raised in cages with wind deflectors that deceived the birds about wind direction. When released in unfamiliar areas, these pigeons showed predictable errors in orientation (Baldaccini et al. 1975, 1978, Fiaschi et al. 1981). Further experiments to determine whether light (including polarized light), sounds, or infrasounds also were involved in this deflector effect have sometimes shown conflicting results but generally have indicated that odors were of primary importance (Ioale and Benvenuti 1983).

Pigeons commonly return home from hundreds of kilometers away (Walcott 1974), so further experiments were conducted to determine whether olfactory cues also are important at these distances (Ioale et al. 1983). Pigeons released more than about 60 km from home showed poor homing orientation unless they had access to olfactory cues during translocation (Ioale et al. 1983). However, some of the pigeons that were prevented from smelling during translocation oriented homeward (Ioale et al. 1983) as was also reported for completely anosmic pigeons (Walraff 1983). This suggests that pigeons in distant unfamiliar areas use nonolfactory stimuli in addition to olfactory stimuli to determine the homeward direction.

The extent to which black bears use olfactory stimuli for homing has not been tested. However, the homeward orientation of bears that were drugged and unconscious for some or all of their translocation (Harger 1970, Rogers in preparation a) suggests that olfaction during translocation may not be necessary for homing in this species. I thank R. R. Buech, L. D. Mech, and C. Walcott for reviewing the manuscript.

LITERATURE CITED

- Alt, G. L. 1977. Home range, annual activity patterns, and movements of black bears in northeastern Pennsylvania. M.S. thesis. Pennsylvania State University, University Park. 67 pp.
- Baker, R. R., J. G. Mather, and J. H. Kennaugh. 1983. Magnetic bones in human sinuses. *Nature* 301:78-80.
- Benvenuti, S., V. Fiaschi, I. Fiore, and F. Papi. 1973a. Homing performances of inexperienced and directionally trained pigeons subjected to olfactory nerve section. *J. Comp. Physiol.* 83:81-92.
- . 1973b. Disturbances of homing behavior in pigeons experimentally induced by olfactory stimuli. *Monitore Zool. Ital.* 7:117-128.
- Delius, J. D., and J. Emmerton. 1978. Sensory mechanisms related to homing in pigeons. Pages 35-41 in K. Schmidt-Koenig and W. T. Keeton, eds. *Animal migration, navigation, and homing*. Springer-Verlag, Berlin-Heidelberg, and New York. 462 pp.
- , R. J. Rerchard, and J. A. Emmerton. 1976. Polarized light discrimination by pigeons and an electroretinographic correlate. *J. Comp. Physiol. Psychol.* 90:560-571.

- Emien, S. T. 1975. Migration, orientation, and navigation. Pages 129-219 in D. S. Farner, J. R. King, and K. C. Parkes, eds. *Avian biology*, vol. V Academic Press, New York, San Francisco, and London. 523 pp.
- Fiaschi, V., N. E. Baldaccini, P. Iola, and F. Papi. 1981. Helicopter observations of homing pigeons with biased orientation because of deflected winds at the home loft. *Monitore Zool. Ital.* 15:139-153.
- Frankel, R. B., R. P. Blakemore, R. S. Wolfe. 1979. Magnetite in freshwater magnetotactic bacteria. *Science* 203:1355-1356.
- Gould, J. L., J. L. Kirschvink, and K. S. Deffeyes. 1978. Bees have magnetic remanence. *Science* 201:1026-1028.
- Griffin, D. R. 1955. Bird navigation. Pages 154-197 in A. Wolfson, ed. *Recent studies in avian biology*. University of Illinois Press, Urbana.
- Harger, E. M. 1970. A study of homing behavior of black bears. M.A. thesis. Northern Michigan University, Marquette. 81 pp.
- Iola, P., and S. Benvenuti. 1983. Pigeon homing: further experiments on shielded lofts. *Comp. Biochem. Physiol.* 76A: 725-731.
- , H. G. Walraff, F. Papi, and A. Foa. 1983. Long-distance releases to determine the spatial range of pigeon navigation. *Comp. Biochem. Physiol.* 76A:733-742.
- Kenyon, K. W., and D. W. Rice. 1958. Homing of Laysan albatrosses. *Condor* 60:3-6.
- Kreithen, M. I. 1978. Sensory mechanisms for animal orientation—can any new ones be discovered? Pages 25-34 in K. Schmidt-Koenig and W. T. Keeton, eds. *Animal migration, navigation, and homing*. Springer-Verlag, Berlin, Heidelberg, and New York. 462 pp.
- Matthews, G. V. T. 1968. *Bird navigation*, second edition. Cambridge University Press, London and New York. 197 pp.
- Mazzeo, R. 1953. Homing of the Manx shearwater. *Auk* 70:200-201.
- Papi, F., L. Fiore, V. Fiaschi, and S. Benvenuti. 1978. An experiment for testing the hypothesis of olfactory navigation of homing pigeons. *J. Comp. Physiol.* 83:93-102.
- . 1984. Homing by black bears and other large mammals. Pages 000-000 in D. S. Maehr and J. R. Brady eds. *Proc. East. Workshop Black Bear Manage, and Res. Vol. 7*. Florida Game and Fresh Water Fish Commission, Gainesville, FL.
- . In preparation a. Navigation by adult black bears.
- . In preparation b. Homing by radio-collared black bears in northeastern Minnesota.
- Southern, W. E. 1959. Homing of purple martins. *Wilson Bull.* 71:254-261.
- . 1968. Experiments on the homing behavior of purple martins. *Living Bird* 7:71-84.
- Walcott, C. 1974. The homing of pigeons. *American Scientist* 62:542-552.
- , J. L. Gould, and J. L. Kirschvink. 1979. Pigeons have magnets. *Science* 205:1027-1029.
- Walraff, H. G. 1983. Relevance of atmospheric odours and geomagnetic field to pigeon navigation: what is the "map" basis? *Comp. Biochem. Physiol.* 76A:643-663.
- Zoeger, J., J. R. Dunn, and M. Fuller. 1981. Magnetic material in the head of the common pacific dolphin. *Science* 213:892-894.